

# Species-specific differences in peroxisome proliferation, catalase, and SOD2 upregulation as well as toxicity in human, mouse, and rat hepatoma cells induced by the explosive and environmental pollutant 2,4,6-trinitrotoluene

Naumenko E., Ahlemeyer B., Baumgart-Vogt E.  
*Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia*

---

## Abstract

© 2016 Wiley Periodicals, Inc. 2,4,6-Trinitrotoluene (TNT) has been widely used as an explosive substance and its toxicity is still of interest as it persisted in polluted areas. TNT is metabolized in hepatocytes which are prone to its toxicity. Since analysis of the human liver or hepatocytes is restricted due to ethical reasons, we investigated the effects of TNT on cell viability, reactive oxygen species (ROS) production, peroxisome proliferation, and antioxidative enzymes in human (HepG2), mouse (Hepa 1-6), and rat (H4IIEC3) hepatoma cell lines. Under control conditions, hepatoma cells of all three species were highly comparable exhibiting identical proliferation rates and distribution of their cell cycle phases. However, we found strong differences in TNT toxicity with the lowest IC 50 values (highest cell death rate) for rat cells, whereas human and mouse cells were three to sevenfold less sensitive. Moreover, a strong decrease in cellular dehydrogenase activity (MTT assay) and increased ROS levels were noted. TNT caused peroxisome proliferation with rat hepatoma cells being most responsive followed by those from mouse and human. Under control conditions, rat cells contained fivefold higher peroxisomal catalase and mitochondrial SOD2 activities and a twofold higher capacity to reduce MTT than human and mouse cells. TNT treatment caused an increase in catalase and SOD2 mRNA and protein levels in human and mouse, but not in rat cells. Similarly, human and mouse cells upregulated SOD2 activity, whereas rat cells failed therein. We conclude that TNT induced oxidative stress, peroxisome proliferation and mitochondrial damage which are highest in rat cells rendering them most susceptible toward TNT. © 2016 Wiley Periodicals, Inc. *Environ Toxicol* 32: 989–1006, 2017.

<http://dx.doi.org/10.1002/tox.22299>

---

## Keywords

catalase, hepatoma cells, human, mouse, peroxisome proliferation, PEX14, rat, SOD2, superoxide radical anion, TNT

## References

- [1] Ahlborg G Jr, Sorsa M. 1988. Mutagenic activity and metabolites in the urine of workers exposed to trinitrotoluene (TNT). *Br J Ind Med* 45:353-358.
- [2] Ahlemeyer B, Neubert I, Kovacs WJ, Baumgart-Vogt E. 2007. Differential expression of peroxisomal matrix and membrane proteins during postnatal development of mouse brain. *J Comp Neurol* 505:1-17.
- [3] Ahlemeyer B, Vogt JF, Michel V, Hahn-Kohlberger P, Baumgart-Vogt E. 2014. Microporation is an efficient method for siRNA-induced knockdown of PEX5 in HepG2 cells: Evaluation of the transfection efficiency, the PEX5 mRNA and protein levels and induction of peroxisomal deficiency. *Histochem Cell Biol* 142:577-591.
- [4] Ammerschläger M, Beigel J, Klein KU, Müller SO. 2004. Characterization of species specificity of peroxisome proliferation in rat and human hepatocytes. *Toxicol Sci* 78:229-240.
- [5] Arlt VM, Hewer A, Sorg BL, Schmeiser HH, Phillips DH, Stiborova M. 2004. 3-aminobenzanthrone, a human metabolite of the environmental pollutant 3-nitrobenzanthrone, forms DNA adducts after metabolic activation by human and rat liver microsomes: Evidence for activation by cytochrome P450 1A1 and P450 1A2. *Chem Res Toxicol* 17:1092-1101.
- [6] Bai J, Rodriguez AM, Melendez JA, Cederbaum AI. 1999. Overexpression of catalase in cytosolic or mitochondrial compartment protects HepG2 cells against oxidative injury. *J Biol Chem* 274:2617-2624.
- [7] Banerjee HN, Verma M, Hou LH, Ashraf M, Dutta SK. 1999. Cytotoxicity of TNT and its metabolites. *J Biol Med* 72:1-4.
- [8] Baumgart E. 1997. Application of in situ hybridization, cytochemical and immunocytochemical techniques for the investigation of peroxisomes. *Histochem Cell Biol* 108:185-210.
- [9] Berthe-Corti L, Jacobi H, Kleihauer S, Witte I. 1998. Cytotoxicity and mutagenicity of a 2,4,6-trinitrotoluene (TNT) and hexogen contaminated soil in *S. typhimurium* and mammalian cells. *Chemosphere* 37:209-218.
- [10] Berridge MV, Herst PM, Tan AS. 2005. Tetrazolium dyes as tools in cell biology: New insights into their cellular reduction. *Biotechnol Annu Rev* 11:127-152.
- [11] Bolt HM, Degen GH, Dorn SB, Plöttner S, Harth V. 2006. Genotoxicity and potential carcinogenicity of 2,4,6-TNT trinitrotoluene: Structural and toxicological considerations. *Rev Environ Health* 21:217-228.
- [12] Borbath I, Leclercq I, Moulin P, Sempoux C, Horsmans Y. 2007. The PPARgamma agonist pioglitazone inhibits early neoplastic occurrence in the rat liver. *Eur J Cancer* 43:1755-1763.
- [13] Bradford MM. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein dye binding. *Anal Biochem* 72:248-254.
- [14] Brocard C, Es-Souni M, Ramirez LC, Latruffe N, Bournot P. 1993. Stimulation of peroxisomal palmitoyl-CoA oxidase activity by ciprofibrate in hepatic cell lines: Comparative studies in Fao, MH1C1 and HepG2 cells. *Biol Cell* 77:37-41.
- [15] Bueding E, Jolliffe N. 1946. Metabolism of trinitrotoluene (TNT) in vitro. *J Pharmacol Exp Ther* 88:300-312.
- [16] Butterworth BE, Earle LL, Strom S, Jirtle R, Michalopoulos G. 1983. Induction of DNA repair in human and rat hepatocytes by 1,6-dinitropyrene. *Mutat Res* 122:73-80.
- [17] Channon HJ, Mills GT, Williams RT. 1944. The metabolism of 2:4:6-trinitrotoluene (alpha-T.N.T.). *Biochem J* 38:70-85.
- [18] Chien CC, Kao CM, Chen DY, Chen SC, Chen CC. 2014. Biotransformation of trinitrotoluene (TNT) by *Pseudomonas* spp. isolated from a TNT-contaminated environment. *Environ Toxicol Chem* 33:1059-1063.
- [19] Cho YS, Lee BU, Kahng HY, Oh KH. 2009. Comparative analysis of 2,4,6-trinitrotoluene (TNT)-induced cellular responses and proteomes in *Pseudomonas* sp. HK-6 in two types of media. *J Microbiol* 4:220-224.
- [20] Colasante C, Chen J, Ahlemeyer B, Baumgart-Vogt E. 2015. Peroxisomes in cardiomyocytes and the peroxisome/proxisome proliferator-activated receptor-loop. *Thromb Haemost* 113:425-436.
- [21] Cornu-Chagnon MC, Dupont H, Edgar A. 1995. Fenofibrate: Metabolism and species differences for peroxisome proliferation in cultured hepatocytes. *Fundam Appl Toxicol* 26:63-74.
- [22] Deng Y, Meyer SA, Guan X, Escalon BL, Ai J, Wilbanks MS, Welti R, Garcia-Reyero N, Perkins EJ. 2011. Analysis of common and specific mechanisms of liver function affected by nitrotoluene compounds. *PLoS One* 6:e14662.
- [23] Elcombe CR, Bell DR, Elias E, Hasmall SC, Plant NJ. 1986. Peroxisome proliferators: Species differences in response of primary hepatocyte cultures. *Ann NY Acad Sci* 804:628-635.
- [24] El-Naa NM, El-Refaei MF, Nasif WA, Abduljawad SH, El-Bairry AI, El-Readi MZ. 2015. In-vivo antioxidant and anti-inflammatory activity of rosiglitazone, a peroxisome proliferator-activated receptor-gamma (PPAR-γ) agonists in animal model of bronchial asthma. *J Pharm Pharmacol* 67:1421-1430.
- [25] Esteve-Nunez A, Caballero A, Ramos JL. 2001. Biological degradation of 2,4,6-trinitrotoluene. *Microbiol Mol Biol Rev* 65:335-352.
- [26] Fahimi HD, Baumgart E, Völkl A. 1993. Ultrastructural aspects of the biogenesis of peroxisomes in rat liver. *Biochimie* 75:201-208.

- [27] Fahimi HD, Beier K, Lindauer M, Schad A, Zhan J, Pill J, Rebel W, Völkl A, Baumgart E. 1996. Zonal heterogeneity of peroxisome proliferation in rat liver. In: Reddy JK, Suga T, Mannaerts GP, Lazarow PB, Subramani S, editors. *Peroxisomes. Biology and Role in Toxicology and Disease*. New York, NY: New York Academy of Sciences. Vol. 804, pp 341–361.
- [28] Fransen M, Nordgren M, Wang B, Apanasets O. 2011. Role of peroxisomes in ROS/RNS metabolism: Implications for human disease. *Biochim Biophys Acta* 1822:1363–1373.
- [29] George SE, Kohan MJ, Warren SH. 1996. Hepatic DNA adducts and production of mutagenic urine in 2,6-dinitrotoluene-treated B6C3F1 male mice. *Cancer Lett* 102:107–111.
- [30] Glass KY, Newsome CR, Tchounwou PB. 2005. Cytotoxicity and expression of c-fos, HSP70, and GADD45/153 proteins in human liver carcinoma (HepG2) cells exposed to dinitrotoluenes. *Int J Environ Res Public Health* 2:355–361.
- [31] Gonzalez FJ, Shah YM. 2008. PPARalpha: Mechanism of species differences and hepatocarcinogenesis of peroxisome proliferators. *Toxicology* 246:2–8.
- [32] Goodsell DS. 2004. Catalase. Molecule of the Month. RCSB Protein Data Bank.
- [33] Grant P, Ahlemeyer B, Karnati S, Berg T, Stelzig I, Nenicu A, Kuchelmeister K, Crane DI, Baumgart-Vogt E. 2013. The biogenesis protein PEX14 is an optimal marker for the identification and localization of peroxisomes in different cell types, tissues, and species in morphological studies. *Histochem Cell Biol* 140:423–442.
- [34] Homma-Takeda S, Hiraku Y, Ohkuma Y, Oikawa S, Murata M, Ogawa K, Iwamuro T, Li S, Sun GF, Kumagai Y, Shimojo N, Kawanishi N. 2002. 2,4,6-trinitrotoluene-induced reproductive toxicity via oxidative DNA damage by its metabolite. *Free Radic Res* 36:555–566.
- [35] Homma-Takeda S, Hiraku Y, Ohkuma Y, Oikawa S, Murata M, Ogawa K, Iwamuro T, Li S, Sun Joo HJ, Yim YH, Jeong PY, Jin YX, Lee JE, Kim H, Jeong SK, Chitwood DJ, Paik YK. 2009. *Caenorhabditis elegans* utilizes dauer pheromone biosynthesis to dispose of toxic peroxisomal fatty acids for cellular homeostasis. *Biochem J* 422:61–71.
- [36] Immenschuh S, Baumgart-Vogt E. 2005. Peroxiredoxins, oxidative stress, and cell proliferation. *Antioxid Redox Signal* 7:768–777.
- [37] Islinger M, Li KW, Seitz J, Völkl A, Lüers GH. 2009. Hitchhiking of Cu/Zn superoxide dismutase to peroxisomes—Evidence for a natural piggyback import mechanism in mammals. *Traffic* 10:1711–1721.
- [38] Karnati S, Baumgart-Vogt E. 2008. Peroxisomes in mouse and human lung: Their involvement in pulmonary lipid metabolism. *Histochem Cell Biol* 130:719–740.
- [39] Kiria-Sakai M, Kumagai Y, Li S, Shimojo N. 1995. Generation of hydrogen peroxide during reduction of 2,4,6-trinitrotoluene and its related compounds. *Sangyo Eiseigaku Zasshi* 41:11–12.
- [40] Kumagai Y, Kikushima M, Nakai Y, Shimojo N, Kunitomo M. 2004. Neuronal nitric oxide synthase (nNOS) catalyses one-electron reduction of 2,4,6-trinitrotoluene, resulting in decreased nitric oxide production and increased nNOS gene expression: Implication for oxidative stress. *Free Rad Biol Med* 37:350–357.
- [41] Lachance B, Robidoux PY, Hawari J, Ampleman G, Thiboutot S, Sunahara GI. 1999. Cytotoxic and genotoxic effects of energetic compounds on bacterial and mammalian cells in vitro. *Mutat Res* 444:25–39.
- [42] Lai DY. 2004. Rodent carcinogenicity of peroxisome proliferators and issues on human relevance. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 22:37–55.
- [43] Miliukiene V, Cenas N. 2008. Cytotoxicity of nitroaromatic explosives and their biodegradation products in mice splenocytes: Implications for their immunotoxicity. *Z Naturforsch* 63:519–525.
- [44] Mukherjee R, Jow L, Noonan D, McDonnell DP. 1994. Human and rat peroxisome proliferator activated receptors (PPARs) demonstrate similar tissue distribution but different responsiveness to PPAR activators. *J Steroid Biochem Mol Biol* 51:157–166.
- [45] Naumenko EA, Naumov AV, Suvorova ES, Gerlach R, Ziganshin AM, Lozhkin AP, Naumova RP. 2008. Participation of oxygen in the bacterial transformation of 2,4,6-trinitrotoluene. *Biochemistry (Moscow)* 73:463–469.
- [46] Naumenko EA, Sibgatullina GV, Mukhitov AR, Rodionov AA, Ilinskaya ON, Naumova RP. 2013. 2,4,6-Trinitrotoluene as a trigger for oxidative stress in *Fagopyrum taticum* callus cells. *Russ J Plant Physiol* 60:404–410.
- [47] O'Brien ML, Spear BT, Glauert HP. 2005. Role of oxidative stress in peroxisome proliferator-mediated carcinogenesis. *Crit Rev Toxicol* 35:61–88.
- [48] Oswal DP, Balanarasimha M, Loyer JK, Bedi S, Soman FL, Rider SD Jr, Hostetler HA. 2013. Divergence between human and murine peroxisome proliferator-activated receptor alpha ligand specificities. *J Lipid Res* 54:2354–2365.
- [49] Reddy JK, Azarnoff DL, Hignite CE. 1980. Hypolipidaemic hepatic peroxisome proliferators form a novel class of chemical carcinogens. *Nature* 283:397–398.

- [50] Rodriguez C, Noe V, Cabrero A, Ciudad CJ, Laguna JC. 2000. Differences in the formation of PPARalpha-RXR/jacoPPRE complexes between responsive and nonresponsive species upon fibrate administration. *Mol Pharmacol* 58:185-193.
- [51] Sarlauskas J, Nemeikaite-Ceniene A, Anusevicius Z, Miseviciene L, Julvez MM, Medina M, Gomez-Moreno C, Cenas N. 2004. Flavoenzyme-catalyzed redox cycling of hydroxylamino- and amino metabolites of 2,4,6-trinitrotoluene: Implications for their cytotoxicity. *Arch Biochem Biophys* 425:184-192.
- [52] Schrader M, Wodopia R, Fahimi HD. 1999. Induction of tubular peroxisomes by UV irradiation and reactive oxygen species in HepG2 cells. *Histochem Cytochem* 47:1141-1148.
- [53] Schrader M, Fahimi HD. 2006. Peroxisomes and oxidative stress. *Biochim Biophys Acta* 1763:1755-1766.
- [54] Stangl H, Kovacs W, Böck P, Kremser K. 1995. Differential induction of peroxisomal enzymes by hypolipidaemics in human (HepG2) and rat (MH1C1) hepatoma cell lines. *Eur J Clin Chem Clin Biochem* 33:775-783.
- [55] Stenuit B, Lamblin G, Cornelis P, Agathos SN. 2012. Aerobic denitration of 2,4,6-trinitrotoluene in the presence of phenazine compounds and reduced pyridine nucleotides. *Environ Sci Technol* 46:10605-10613.
- [56] Stier H, Fahimi HD, Van Veldhoven P, Mannaerts GP, Völkl A, Baumgart E. 1998. Maturation of peroxisomes in differentiating human hepatoblastoma cells (HepG2): Possible involvement of the peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ). *Differentiation* 64:55-66.
- [57] Taub J, Lau JF, Ma C, Hahn JH, Hoque R, Rothblatt J, Chalfie M. 1999. A cytosolic catalase is needed to extend adult lifespan in *C. elegans* daf-C and clk-1 mutants. *Nature* 399:162-166.
- [58] Tchounwou PB, Wilson BA, Ishaque AB, Schneider J. 2001. Transcriptional activation of stress genes and cytotoxicity in human liver carcinoma cells (HepG2) exposed to 2,4,6-trinitrotoluene, 2,4-dinitrotoluene, and 2,6-dinitrotoluene. *Environ Toxicol* 16:209-216.
- [59] Thiele S, Fernandes E, Bollag JM. 2002. Enzymatic transformation and binding of labeled 2,4,6-trinitrotoluene to humic substances during an anaerobic/aerobic incubation. *J Environ Qual* 31:437-444.
- [60] Tugwood JD, Aldridge TC, Lambe KG, Macdonald N, Woodyatt NJ. 1996. Peroxisome proliferator-activated receptors: Structures and function. *Ann NY Acad Sci* 804:252-265.
- [61] Woodyatt NJ, Lambe KG, Myers KA, Tugwood JD, Roberts RA. 1999. The peroxisome proliferator (PP) response element upstream of the human acyl CoA oxidase gene is inactive among a sample human population: Significance for species differences in response to PPs. *Carcinogenesis* 20:369-372.
- [62] Xiao Y, Karnati S, Qian G, Nenicu A, Fan W, Tchatalbachev S, Höland A, Hossain H, Guillou F, Lüers GH, Baumgart-Vogt E. 2012. Cre-mediated stress affects sirtuin expression levels, peroxisome biogenesis and metabolism, antioxidant and proinflammatory signaling pathways. *PLoS One* 7:e41097.
- [63] Yu J, Shen B, Chu ES, Teoh N, Cheung KF, Wu CW. 2010. Inhibitory role of peroxisome proliferator-activated receptor gamma in hepatocarcinogenesis in mice and in vitro. *Hepatology* 51:2008-2019.
- [64] Zaripov SA, Naumov AV, Suvorova ES, Garusov AV, Naumova RP. 2004. Initial stages of 2,4,6-trinitrotoluene transformation by microorganisms. *Microbiology* 73:398-403.
- [65] Zhang XC, Hu JP. 2008. FISSON1A and FISSON1B proteins mediate the fission of peroxisomes and mitochondria in *Arabidopsis*. *Mol Plant* 1:1036-1047.
- [66] Ziganshin AM, Gerlach R, Borch T, Naumov AV, Naumova RP. 2007. Production of eight different hydride complexes and nitrite release from 2,4,6-trinitrotoluene by *Yarrowia lipolytica*. *Appl Environ Microbiol* 73:7898-7905.
- [67] Ziganshin AM, Naumov AV, Suvorova ES, Naumenko EA, Naumova RP. 2010. Hydride-mediated reduction of 2,4,6-trinitrotoluene by yeasts as the way to its deep degradation. *Mikrobiologika* 76:766-773.
- [68] Ziganshin AM, Ziganshina EE, Byrne J, Gerlach R, Struve E, Biktagirov T, Rodionov A, Kappler A. 2015. Fe III mineral reduction followed by partial dissolution and reactive oxygen species generation during 2,4,6-trinitrotoluene transformation by aerobic yeast *Yarrowia lipolytica*. *AMB Express* 5:8-20.